


- 1 The table below gives data from tests of a full-size bungee jump.

Jumper Weight (in pounds)	100	125	150	175	200
Stretched Cord Length (in feet)	50	55	60	65	70

- a. Which variable does it make sense to consider independent and which dependent?
- b. Plot the given data on a coordinate graph.



- c. Use the pattern in the table or the graph to estimate the stretched cord length for jumpers who weigh:
- i. 85 pounds                      ii. 135 pounds                      iii. 225 pounds

d. Would it make sense to connect the points on your data plot? Explain your reasoning.


e. Describe the overall pattern relating *stretched cord length*  $L$  to *jumper weight*  $w$ .

f. The technician who did the tests suggested that the pattern could be summarized with a symbolic rule  $L = 30 + 0.2w$ . Does that rule give estimates of stretched cord length that match the experimental data? Explain.

**17** When there appears to be a relationship between values of two variables, how do you decide which should be considered the *independent variable* and which should be considered the *dependent variable*?

**7** In 2012, the Olympic record in the women’s 100-meter freestyle swim race was 53.00 seconds. It was set by Ranomi Kromowidjojo from the Netherlands. She swam at an average speed of  $100 \div 53 \approx 1.88$  meters per second.

- a. Make a table and a graph showing the way *average speed* for the 100-meter race changes as *time* increases from 40 seconds to 120 seconds (2 minutes) in steps of 10 seconds.

- b. Describe the pattern of change shown in your table and graph.

c. Write a rule showing how to calculate *average speed*  $s$  for any *race time*  $t$ .

d. Which change in *race time* will cause the greatest change in *average speed*:  
an increase from 50 to 60 seconds or an increase from 110 to 120 seconds?  
Explain how your answer is illustrated in the shape of your graph.